arXiv:cond-mat/0206106v1 7 Jun 2002

Comment on "Spin Dynamics of the 2D Spin $\frac{1}{2}$ Quantum Antiferromagnet Copper Deuteroformate Tetradeuterate (CFTD)" :

In a recent Letter [1] Rønnow *et al.* presented a neutron scattering study of the magnetic excitations of the antiferromagnet $\mathrm{Cu(DCOO)_2 \cdot 4D_2O}$ for temperatures up to $T \approx J/2$, where $J \approx 73K$ is the nearest neighbor exchange energy. Due to the smallness of the interlayer-coupling and anisotropy-energies, this system is an almost ideal realization of a square lattice nearest neighbor spin $S = \frac{1}{2}$ quantum Heisenberg antiferromagnet. Rønnow *et al.* paid particular attention to the temperature-dependence of the dispersion and the damping of magnons with wave-vectors $\mathbf{k}$ close to the boundary of the magnetic Brillouin zone. They observed a "surprisingly good" agreement of the data for the temperature-dependence of the magnon-damping rate Γ with the formula

$$\Gamma_{\rm long}(T) \approx v_s(T)/\xi(T) \; , \tag{1}$$

which expresses the damping of *long-wavelength* magnons in terms of the spin-wave velocity $v_s(T)$ and the antiferromagnetic correlation length $\xi(T)$.

In this Comment we point out that the agreement of the experimental data with Eq. (1) is accidental, because a comparison with Eq. (1) is only meaningful for *long-wavelength* magnons [2], whereas the experimental data give the damping $\Gamma_{\rm short}(T)$ of *short-wavelength* magnons, with wave-vectors close to the boundary of the magnetic Brillouin zone. The experimentally relevant damping rate of short-wavelength magnons for spin S Heisenberg antiferromagnets has been calculated in Ref. [3]. For wave-vectors in the regime $|\mathbf{k}| \gtrsim (Ta/v_s)^{1/3}(2\pi/a)$ (which includes the zone boundary) the damping is [3,4]

$$\frac{\Gamma_{\rm short}(T)}{J} = \frac{\pi}{24}\frac{Z(|\mathbf{v_k}|)}{S^4}\left(\frac{T}{J}\right)^3 \left[1 + O\left(\frac{1}{2S}, \frac{T}{4SJ}\right)\right] \; , \tag{2}$$

where the numerical factor $Z(|\mathbf{v_k}|)$ depends on the length of the dimensionless magnon-velocity $\mathbf{v_k} = \sqrt{2}\nabla_{\mathbf{k}}\epsilon_{\mathbf{k}}$ [5]. From the numerical evaluation of $Z(|\mathbf{v_k}|)$ [3] it is easy to see that $Z(|\mathbf{v_k}|) \approx Z(0) = 1$ for magnons with wave-vectors in the vicinity of the zone boundary, where $|\mathbf{v_k}| \ll 1$. Note that Eq. (2) contains no adjustable parameters.

Taking into account that higher orders in $1/S$ and in $T/(SJ)$ have been neglected in Eq. (2), the agreement with the experimental data [1] is excellent, see Fig. 1. Note that Eq. (2) is based on the evaluation of Fermi's golden rule, using for the relevant matrix elements the bare vertices describing magnon-magnon scattering within the Dyson-Maleev transformation [3]. The fact that the resulting damping rate is in quantitative agreement with the experiment indicates that the effective interaction between antiferromagnetic magnons at short wavelengths is weak and hence can be treated perturbatively, without taking vertex corrections into account [6]. On the other hand, at long wavelengths the effective magnon-magnon interaction becomes strong and prohibits the propagation of stable magnons over length scales exceeding the correlation length $\xi(T)$. This is the physics behind the estimate (1) for the magnon damping rate at long wavelengths, which has nothing to do with the short-wavelength damping due to weak magnon-magnon scattering given in Eq. (2).

This work was supported by the DFG via Forschergruppe FOR 412, Project No. KO 1442/5-1.

---

[1] H. M. Rønnow *et al.* , Phys. Rev. Lett. **87**, 037202 (2001).

[2] More precisely, Eq. (1) can only be correct for $|\mathbf{k}| \ll 2\pi/\xi$. Magnon damping in the regime $2\pi/\xi \lesssim |\mathbf{k}| \ll 2\pi/a$ (where a is the lattice spacing) has been discussed by S. Tyč and B. I. Halperin, Phys. Rev. B **42**, 2096 (1990).

[3] P. Kopietz, Phys. Rev. B **41**, 9228 (1990).

[4] We use units where $\hbar = k_B = 1$ and define $\Gamma \equiv {\rm Im}\Sigma(\mathbf{k}, E_{\mathbf{k}} - i0)$, which differs from the definition used in Ref. [3] by a factor of two, but agrees with Ref. [1].

[5] Here $\epsilon_{\mathbf{k}} = (1 - \gamma_{\mathbf{k}}^2)^{1/2}$, where $\gamma_{\mathbf{k}} = \frac{1}{2}[\cos(k_x a) + \cos(k_y a)]$.

[6] P. Kopietz and S. Chakravarty, Phys. Rev. B **56**, 3338 (1997).

Peter Kopietz and Ivan Spremo
Institut für Theoretische Physik, Universität Frankfurt,
Robert-Mayer-Strasse 8, 60054 Frankfurt, Germany

Received: August 16, 2001
PACS numbers: 75.10.Jm, 05.70.Jk, 75.50.Ee



FIG. 1. Solid line: temperature-dependence of the damping of the zone-boundary magnons, see Eq. (2) for $S = 1/2$. The circles are the experimental data from Fig. 3 of Ref. [1].